



# Prevalence and Microbiological Profile of Catheter Associated Urinary Tract Infections: A Survey in Secondary Care Hospital in Gaza Strip

Aymen Elsous<sup>1,2\*</sup>, Mahmoud Ouda<sup>2</sup>, Samah Mohsen<sup>2</sup>, Mohammed Al-Shaikh<sup>2</sup>, Siham Mokayad<sup>2</sup>

<sup>1</sup> Department of Health Management and Economics, School of Public Health, International Campus, Tehran University of Medical Sciences, Tehran, Iran. <sup>2</sup> Quality Improvement and Infection Control Office, Shifa Medical Complex, Ministry of Health, Gaza Strip, Palestine.

First Published online June 30, 2016

## Abstract

**Background and Objectives:** Healthcare-associated infections is a major health care concern posing potentially serious negative impact on patient safety and outcome. In this paper, we report this microbiological profile and the prevalence of catheter associated urinary tract infections (CAUTIs).

**Methods:** This was a prospective observational study for 60 inpatients from 8 wards among patients having indwelling catheter for at least 48 hours. Urine cultures were taken in initial phase within 24 hours of admission, at 48 hours and in day 5 of catheterization.

**Findings:** The rate of CAUTIs was 16.7% after 48 hours and 28% (7/25) after 5 days from insertion the indwelling catheter. *Candida* spp. was the most common cause of CAUTIs (29.4%), followed by *Escherichia coli* (23.5%), *Streptococcus* spp. and *Klebsiella* spp. (17.6%) and last *Staphylococcus* spp. (11.7%). The coronary care unit (CCU) showed the highest prevalence of CAUTIs (57.1%), followed by Oncology (42.8%), and Internal Medicine (33.3%). Females showed a significantly higher rate of acquired urinary tract infections (UTIs) compared with males ( $P < .05$ ).

**Conclusions:** The risk of CAUTIs was found to be relatively high and increase by duration of catheterization. These results recommend minimally usage of catheter in medical practice in terms of both frequency and time, particularly for female patients and in CCU ward. Identification of the microbiological profile of the CAUTIs would help efficient treatment of the infected patients.

**Keywords:** Catheter associated urinary tract infections, Prevalence, Hospital, Microbiological profile

## Background and Objectives

People who seek healthcare are unable to tolerate the additional burden of acquiring a new infection. Healthcare associated infections (HAIs) are a major concern for healthcare management and frontlines to maximize patient safety. HAI is global health problem because of their seriousness, burden and costs to healthcare system. They also impose serious problems to safety of patients.<sup>1</sup> In developing countries, the prevalence and incidence of HAIs varies between 5.7 to 19.1, and 1.7 to 23.6 per 100 patients respectively.<sup>2</sup> Studies from European hospitals

indicates that 1 in every 18 patients has HAIs.<sup>3</sup> In the United States 1.7 million suffer from HAIs and 100 000 dies every year as result of these infections.<sup>4</sup>

Urinary catheterization occurs in 15%-25% of all hospitalized patients and urinary tract infections (UTIs) caused by urinary catheter are the most common type of HAIs,<sup>5</sup> representing 30% to 40% of all HAIs in some regions of the world.<sup>6-9</sup> Over 80% of catheter associated urinary tract infections (CAUTIs) are linked to insertion of indwelling catheter.<sup>10</sup> The rate of these infections can be reduced through effective infection prevention and control program (ICP).<sup>3,11</sup> It is estimated that strict infection and prevention program in has resulted in 15%-69% reduction in infection and prevention of 9000 deaths from CAUTIs.<sup>12,13</sup>

In Palestine, where ICP is at the beginning and starting

\*Corresponding Author: Aymen Mohammed Elsous, Quality Improvement and Infection Control Office, Shifa medical complex, Ministry of Health, Gaza Strip, Palestine. Tel: +970598926886, Email: aymenelsous65@gmail.com

to evolve, no precise or even estimated data are available on HAIs. Having knowledge and accurate data on HAIs will enable hospitals to develop their strategies and preventive measures to reduce the incidence and complications of HAIs including CAUTIs. The aim of this study was (1) to determine the prevalence rate of CAUTIs, (2) to determine the profile of causative pathogens and (3) risk factors associated with CAUTIs.

## Methods

### Study Design and Setting

This prospective study was carried out in Shifa medical complex which situated in Gaza Strip, Palestine. This complex is the oldest and largest healthcare setting with 535 beds and includes 3 hospitals: surgical, internal medicine, and Gyn/Obst hospitals. Samples were taken from patients of five wards which were selected purposefully based on the frequency of catheter insertion, including internal medicine, intensive care unit (ICU), cardiac, burns and orthopedics wards.

### Data Collection and Sampling

Data on patients' age, sex, disease, and type and place of catheterization, and hospitalization ward were obtained from patient's medical records. Patients who used condoms, supra-pubic catheter, and percutaneous nephrostomy tube and those who used catheter prior to admission to the selected wards were excluded. The plan was to take 3 urine samples from each patient: first, within 12 hours after insertion; second, 48 hours from admission, and third, in day 5 of admission. The urine samples were taken from the bifurcation of Foley's catheter with aseptic technique and put in a sterile container. The samples were immediately taken to microbiology lab for culture preparation.

### Culturing and Identification of Microorganism

A plate with agar was used for culturing the urine sample and results were read after 24 hours incubation in ovum at 37°C. Identification of microorganisms was carried out principally based on their morphology. However, in case of difficulty in identification, analytical profile index (API) technique was used to identify the microorganisms. A loop of 1 micron or 10 microns was used to pick up a drop of urine. The sample was then inoculated into blood agar media. Using the same loop, a second drop of urine was taken and then was inoculated into MacConkey agar media. Both plates were incubated for 24 hours at 37°C. Gram (+) bacteria and *Candida* spp. were identified based on wet mount approach according to which 1 colony with 1 drop of normal saline is placed on slide for direct microscopic examination. Gram (-) bacteria were identified using API

20E (Analytical profile index for enterobacteriaceae 20 test) ID Kit (bioMérieux, France).

### Case Definition

According to the Centers for Disease Control and Prevention (CDC) definition,<sup>14</sup> CAUTI is defined as: "A UTI where an indwelling urinary catheter was in place for > 2 calendar days on the date of event, with the day of device placement being day 1, and an indwelling urinary catheter was in place on the date of event or the day before. If an indwelling urinary catheter was in place for > 2 calendar days and then removed, the UTI criteria must be fully met on the day of discontinuation or the next day."

### Statistical Analysis

Data were analyzed using descriptive statistical methods. SPSS version 20 package was used for data analysis.

### Ethical Consideration

The institutional review board of Shifa medical complex approved the study protocol in accordance with Helsinki Declaration. All patients or their families were briefed about the study purpose and their consent was obtained.

## Results

### Characteristics of the Patients

A total of 128 patients were enrolled from which 213 urine cultures were obtained and analyzed. After initial analysis of samples, samples from 68 patients were excluded because the first urine cultures were positive (asymptomatic bacteriuria) or the patient was discharged before taking the second urine sample. Table 1 presents the characteristics of participants. Of the total enrolled patients, 60% were female and 40% were male. The mean age was 52.8 ± 19.50 years. A majority of cases (41.7%) were hospitalized in ICU and most catheters insertion was taken place in the emergency department (71.7%).

### Prevalence of CAUTIs

The overall prevalence of CAUTIs was 28.3% (17/60). The prevalence was 16.7% (10/60) and 28% (7/25) after 48 hours and 5 days of insertion the indwelling catheter, respectively (Table 2). The coronary care unit (CCU) showed the highest prevalence of CAUTIs (57.1%), followed by internal medicine (oncology) (42.8%), and internal medicine (33.3%) wards.

### Place of CAUTIs

Table 3 shows the distribution of isolated microorganisms based on hospital wards and departments. While 71.6% (43/60) of catheters were inserted in the department of

**Table 1.** Patients' Characteristics

Variables	Total (n = 60)	Male (n = 24)	Female (n = 36)
Age (median ± SD)			
≤ 35 years	20.0 ± 3.89	22.0 ± 3.34	15.0 ± 1.16
36-60 years	55.0 ± 6.87	53.0 ± 7.01	57.5 ± 6.46
>60 years	71.0 ± 7.78	78.0 ± 8.35	70.5 ± 7.52
Gender (%)			
Male	40		
Female	60		
Ward specialty, n (%)			
ICU	25 (41.7)	16 (66.7)	9 (25.0)
Cardiac	7 (11.7)	4 (16.7)	3 (8.3)
Orthopedic	10 (16.7)	1 (1.7)	9 (15)
Internal medicine	18 (30)	3 (5)	15 (25)
Place of insertion			
Emergency	16 (71.7)	7 (29.2)	9 (25.0)
Department	43 (26.7)	17 (70.8)	26 (72.2)
Theatre	1 (1.7)	0 (0.0)	1 (2.8)

Abbreviation: SD, standard deviation.

**Table 2.** Distribution of Causative Agents

Organisms	Over all (n = 60)	48 hours (n = 60)		Day 5 (n = 25)	
	%	n	%	n	%
<i>Staphylococcus</i> spp.	11.7	1	10	1	14.3
<i>Streptococcus</i> spp.	17.6	2	20	1	14.3
<i>E. coli</i>	23.5	2	20	2	28.5
<i>Klebsiella</i> spp.	17.6	3	30	0	-
<i>Candida</i> spp.	29.	2	20	3	42.8

**Table 3.** Distribution of Microorganisms Based on Hospital Wards

Hospital Wards	48 hours (n = 60)		Day 5 (n = 25)	
	n	%	n	%
ICU	3	5.0	1	4.0
CCU	3	5.0	1	4.0
Male orthopedic	0	0.0	0	0.0
Female orthopedic	2	3.3	1	4.0
Male internal medicine	0	0.0	1	4.0
Female internal medicine	0	0.0	1	4.0
Internal medicine (oncology)	2	3.3	2	8.0

admission, 26.7% (16/60) were inserted in emergency department, and only 1 case was inserted in theatre. The prevalence of CAUTIs was higher in wards (27.9%) compared with emergency department (23.5%).

### Cause of CAUTIs

Of all CAUTIs, 7 (41.1%) were due to gram-negative bacteria, 5 (29.4%) were due to *Candida* spp. and 5 (29.4%) were due to gram-positive bacteria. The most frequently isolated causative agent was *Candida* spp. (29.4%), followed by *Escherichia coli* (23.5%), *Klebsiella* spp. and *Streptococcus*

spp. (17.6%), and *Staphylococcus* spp. (11.7%).

### Discussion

In this study, the overall rate of CAUTIs was 28.3%. For comparison, the rate of CAUTIs has been reported to be 3.1%,<sup>16</sup> 10.6%-12.6%,<sup>17</sup> and 10.7%<sup>18</sup> from studies conducted in Egypt, Britain, and Turkey, respectively. Other studies report the risk of bacteriuria with catheterization to be 3%-10% which grows to 100% after 30 days.<sup>19,20</sup>

In this study, CAUTIs were found to be higher among females and long hospitalized patients with catheter, and in CCU which is consistent with previous studies.<sup>18,22-25</sup> However, unlike the study of Mladenovi et al no difference in rate of CAUTIs was observed among patients of different age group.<sup>21</sup>

We identified *Candida* spp. as the most frequently occurring microbe in CAUTIs, followed by *E. coli*, *Klebsiella* spp, *Streptococcus* spp. and *Staphylococcus* spp. A previous study from Serbia reported *Candida* spp, *Pseudomonas aeruginosa*, and *Klebsiella* spp. as the most commonly observed organisms in CAUTIs.<sup>21</sup> Another report from Singapore identified *Candida* spp. as most common cause of CAUTIs.<sup>26</sup> Laupland et al<sup>27</sup> showed *E. coli* and *Candida* spp. were dominant cause of CAUTIs followed by *Enterococcus* spp. and *P. aeruginosa*.

### Conclusions

The risk of CAUTIs was found to be relatively high and increase by duration of catheterization. These results recommend minimally usage of catheter in medical practice in terms of both frequency and time, particularly for female patients and in CCU ward. Identification of the microbiological profile of the CAUTIs would help efficient treatment of the infected patients.

### Abbreviations

(HAI): healthcare acquired infection; (CAUTI): catheter associated urinary tract infection; (CCU): coronary care unit; (ICP): infection control program; (UTI): urinary tract infection.

### Competing Interests

The authors declare no competing interests.

### Authors' Contributions

AME, MAO, SM, SM, MS jointly designed the study. AME, SM, MS, SM were involved in data collection and analysis. AME, MAO made the major contribution towards revision of the manuscript. All authors read and approved the final manuscript.

## Acknowledgments

The authors are grateful to managing board, head nurses, and the nursing staff of the Shifa hospital for providing facilitation for conduction of this study. We also thank the head of laboratory department of Shifa hospital, Mr. Kayed Ahmad and the staff of Microbiology Unit, Mr. Nahed Abd El-Latif, Mrs. Hala Abdo and Mr. Mahmoud Al Jaro for their kind support. We also appreciate Mr. Abd Al-Rahman Hamad for helping us in analysis of the data.

## References

1. Umscheid CA, Mitchell MD, Doshi JA, Agarwal R, Williams K, Brennan PJ. Estimating the proportion of healthcare-associated infections that are reasonably preventable and the related mortality and costs. *Infect Control Hosp Epidemiol.* 2011;32(2):101-114. doi:10.1086/657912.
2. Allegranzi B, Nejad SB, Combescure C, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet* 2011;377(9761):228-241. doi:10.1016/S0140-6736(10)61458-4.
3. European Centre for Disease Prevention and Control (ECDC). Point prevalence survey of healthcare-associated infections and antimicrobial use in European acute care hospitals 2011–2012. Stockholm: ECDC; 2013.
4. Klompas M, Yokoe D. Automated surveillance of healthcare-associated infections. *Clin Infect Dis.* 2009;48(1):1268–1275.
5. Fakhri MG, Shemes SP, Pena ME, et al. Urinary catheter in the emergency department; very elderly women are at high risk for unnecessary utilization. *Am J Infect Control.* 2010;38(9):683.
6. Stamm WE. Catheter-associated urinary tract infections: epidemiology, pathogenesis and prevention. *Am J Med.* 1991;91(3B):65S-71S.
7. Plowman R, Graves N, Griffin M, et al. The socio-economic burden of hospital acquired infection. *Euro Surveill.* 2000;5(4):49-50.
8. Ward TT, Jones SR. Genitourinary tract infections. In: Reese RE, Betts RF, eds. *A Practical Approach to Infectious Diseases.* 4th ed. Boston: Little, Brown, and Company; 1996:472-518.
9. Vincent J, Bihari D, Suter P, et al. The prevalence of nosocomial infections in intensive care units in Europe. Results of the European Prevalence of Infection in Intensive Care (EPIC) Study. *JAMA.* 1995;274(8):639-644.
10. Anderson DJ, Kirkland KB, Kaye KS, et al. Under resourced hospital infection control and prevention programs: Penny wise, pound foolish? *Infect Control Hosp Epidemiol* 2007;28(7):767-773. Doi:10.1086/518518
11. National Health and Medical Research Council. Australian Guidelines for the Prevention and Control of Infection in Healthcare. [http://www.nhmrc.gov.au/\\_files\\_nhmrc/publications/attachments/cd33\\_complete.pdf](http://www.nhmrc.gov.au/_files_nhmrc/publications/attachments/cd33_complete.pdf). Published 2010.
12. Smyth ETM, McIlvenny G, Enstone JE, et al. Four Country Healthcare Associated Infection Prevalence Survey 2006: overview of the results. *J Hosp Infect.* 2008;69(3):230-248.
13. Umscheid C, Mitchell M, Agarwal R, et al. Mortality From Reasonably-Preventable Hospital Acquired Infections. A Preventable Epidemic. Washington, DC; 2008.
14. Catheter-Associated Urinary Tract Infection (CAUTI) Event. CDC website. <http://www.cdc.gov/nhsn/PDFs/pscManual/7pscCAUTIcurrent.pdf>. Accessed January 2014.
15. Maki DG, Tambyah PA. Engineering out the risk for infection with urinary catheters. *Emerg Infect Dis.* 2001;7(2):342-347.
16. El-Kholy A, Saied T, Gaber M, et al. Device-associated nosocomial infection rates in intensive care units at Cairo University hospitals: first step toward initiating surveillance programs in a resource-limited country. *Am J Infect Control.* 2012;40(6):e216-e220. doi:10.1016/j.ajic.2011.12.010.
17. Pickard R, Lam T, MacLennan G, et al. Types of urethral catheter for reducing symptomatic urinary tract infections in hospitalized adults requiring short-term catheterization: multicenter randomized controlled trial and economic evaluation of antimicrobial- and antiseptic-impregnated urethral catheters (the CATHETER trial) *Health Technol Assess.* 2012;16(47):1-197.
18. Temiz E, Piskin N, Aydemir H, et al. Factors associated with catheter-associated urinary tract infections and the effects of other concomitant nosocomial infections in intensive care units. *Scand J Infect Dis.* 2012;44(5):344-349. doi: 10.3109/00365548.2011.639031.
19. Saint S, Lipsky BA, Gould SD. Indwelling urinary catheters: a one-point restraint. *Ann Intern Med.* 2002;137(2):125–127.
20. Warren JW, Tenney JH, Hoopes JM, Muncie HL, Anthony WC. A prospective microbiologic study of bacteriuria in patients with chronic indwelling urethral catheters. *J Infect Dis.* 1982;146(6):719-723.
21. Mladenovi J, Veljovi M, Udovi I, et al. Catheter-associated urinary tract infection in a surgical intensive care unit. *Vojnosanit Pregl.* 2015;72(10):883-888.
22. van der Kooij TII, de Boer AS, Manniën J, et al. Incidence and risk factors of device-associated infections and associated mortality at the intensive care in the Dutch surveillance system. *Intensive Care Med.* 2007;33(2):271–278.

23. Bochicchio GV, Joshi M, Shih D, Bochicchio K, Tracy K, Scalea TM. Reclassification of urinary tract infections in critically ill trauma patients: a time-dependent analysis. *Surg Infect*. 2003;4(4):379-385.
24. Leone M, Albanese J, Garnier F, et al. Risk factors of nosocomial catheter-associated urinary tract infection in a polyvalent intensive care unit. *Intensive Care Med*. 2003;29(6):929-932.
25. Foxman B. Epidemiology of urinary tract infections: Incidence, morbidity, and economic costs. *Am J Med*. 2002;113(Suppl 1A):5-13.
26. Tay MK, Lee JY, Wee IY, Oh HM. Evaluation of intensive care unit-acquired urinary tract infections in Singapore. *Ann Acad Med Singap*. 2010;39(6):460-465.
27. Laupland KB, Bagshaw SM, Gregson DB, Kirkpatrick AW, Ross T, Church DL. Intensive care unit-acquired urinary tract infections in a regional critical care system. *Crit Care*. 2005;9(2):R60-R65.

**Please cite this article as:**

Elsous A, Ouda M, Mohsen S, Al-Shaikh M, Mokayad S. Prevalence and microbiological profile of catheter associated urinary tract infections: a survey in secondary care hospital in Gaza Strip. *Int J Hosp Res*. 2016;5(2):69-73. doi:10.15171/ijhr.2016.13.